

# Obscured AGN in WISE

Roberto J. Assef

*Universidad Diego Portales, Chile*

In collaboration with Daniel Stern (JPL), Peter Eisenhardt (JPL), Jingwen Wu (UCLA), Chao-Wei Tsai (JPL), Andrew Blain (Leicester), Christopher S. Kochanek (OSU), Dominika Wylezalek (JHU) and the WISE ExGal Team



# Outline

- “Normal” Obscured AGN with WISE
  - Stern et al.(2012, ApJ, 753, 30)
  - Assef et al.(2013, ApJ, 772, 26)
- Hot Dust Obscured Galaxies - Hot DOGs
  - Eisenhardt et al. (2012, ApJ, 755, 173)
  - Wu et al.(2012, ApJ, 756, 96), Wu et al.(2014, ApJ, 793, 8)
  - Stern et al.(2014, ApJ, 794, 102)
  - Jones et al.(2014, MNRAS, 443, 146)
  - Assef et al.(submitted, arXiv:1408.1092)
  - Tsai et al. (submitted, arXiv:1410.1751)

# Why Care About Obscured AGN?

- AGN likely play a fundamental role in driving galaxy evolution.
- Relations between host and black hole imply a strong degree of co-evolution.
- One of the most expected phases of massive galaxies is an obscured phase where AGN feedback starts shutting-down star-formation.

See Nadia's Talk!



Normal AGN in WISE

# The Role of WISE in Finding AGN


Wright et al. (2010)

The mid-IR is great for identifying AGN


- Particularly at 3-5 microns
- Hot dust from the Torus is particularly bright and blue
- Most stars are red (Rayleigh-Jeans)

We have known this since Spitzer

- Lacy et al. (2004, 2007), Stern et al.(2005), Alonso-Herrero et al. (2008), Donley et al.(2008, 2012), Assef et al.(2010)



# WISE AGN Selection



Most IRAC AGN have


$$W1 - W2 > 0.8$$

- 95% reliability in COSMOS
- 75% completeness in COSMOS

*Stern et al. (2012, ApJ, 753, 30)*

See *Assef et al. (2013, ApJ, 772, 26)* for a more universal AGN selection based on W1 and W2.

# WISE AGN Selection



Most IRAC AGN have

$$W1 - W2 > 0.8$$


- 95% reliability in COSMOS
- 75% completeness in COSMOS

*Stern et al. (2012, ApJ, 753, 30)*

See *Assef et al. (2013, ApJ, 772, 26)* for a more universal AGN selection based on W1 and W2.


# Magnitude De

- W1 and W2 alone a depths
  - Different selection c completeness
  - Our 90% reliability c
    - 6x nominal surfa
    - Increase is prima
  - W3 is great to includ
    - Jarrett et al. (201




# Number of AGN vs. Accretion Disk Obscuration in Bootes

Assef et al. (2013)



# Evidence for a Receding Torus?



# Hot Dust Obscured Galaxies (Hot DOGs)

*Assef et al. (submitted, arXiv:1408.1092)*

# WISE – Most Luminous Galaxies

- One goal of WISE is to identify the most luminous galaxies in the universe.
- These galaxies should be **very red and bright in the mid-IR (SF/AGN)** but most likely **very rare**.


Combination of depth + area makes WISE ideal for this.

- As expected, WISE has been very successful at it with the discovery of **Hot Dust Obscured Galaxies (Hot DOGs)**.

As you have already heard from Chao-Wei Tsai's talk.

Observed-Frame  $\lambda(\mu\text{m})$

1 10 100 1000



First object studied in detail (*Eisenhardt et al. 2012, 755, 173*)

Best fit SED model has


- AGN with  $A_V \sim 50$  mag
- Starburst with  $\sim 300 M_{\text{sun}}/\text{yr}$  SFR
- $L_{\text{Bol}} = 9 \times 10^{13} L_\odot$

*Adapted from Eisenhardt et al.(2012)*


# Hot DOGs are at High-z and are Hyper-Luminous

*Assef et al. (submitted)*

- All at  $z > 1$
- Low contamination rate of selection by low-z interlopers
- Highest at  $z=4.6$   
Will back to this one later on
- All have  $L_{\text{IR}} > 10^{13} L_{\odot}$   
10% have  $L_{\text{IR}} > 10^{14} L_{\odot}$   
*Wu et al. (2012)*  
*Tsai et al. (2015)*




# Remarkably Similar SEDs



*Adapted from Wu et al. (2012, ApJ, 756, 96)*


# Powered by Very Luminous AGN



- 52 Hot DOGs with Spitzer follow-up Spec-z  $W4 < 7.2$  mag
- Similar SED modeling as for W1814+342
- Significantly brighter than  $L^*$
- Among the most luminous AGN

# Maximal Stellar Mass

*Assef et al. (submitted)*



Too few photometric bands to constrain M/L

Upper bound on M/L from

Maximally old stellar population

Bruzual & Charlotte (2003) for low TP-AGB contribution


Bottom-heavy IMF from Conroy & van Dokkum (2013)

# Hot DOGs & Local BH-Host Relations

Hot DOGs have very luminous AGN in not so massive galaxies.

Two possibilities

1. BHs much more massive than expected from local relations
2. Super-Eddington luminosities



*Assef et al. (submitted)*


# Hot DOGs & Local BH-Host Relations

Assuming the local  $M_{\text{BH}} - M_{\text{sph}}$  relation

Hot DOGs have very luminous AGN in not so massive galaxies.


Two possibilities

1. BHs much more massive than expected from local relations
2. Super-Eddington luminosities



*Assef et al. (submitted)*

# Similar Space Density as Type 1 QSOs




- Although rare, Hot DOGs seem to be as common as Type 1 QSOs.  
Type 1 QLF from SDSS/BOSS (Ross et al. 2013)

- Half of the most luminous AGN are obscured?

- No clear redshift evolution  
Sample may be too small

*Assef et al. (submitted)*

# Hot DOGs Live in Dense Environments




- Based on the projected density of galaxies in IRAC within 1 arcmin.
- More dense than the field (SpUDS)
- Similarly dense to proto-clusters at  $z \sim 2$  (CARLA)


*Assef et al. (submitted)*

# W2246-0526, The Most Luminous Hot Dog

*Diaz-Santos et al. (in prep.)*



# W2246-0526, The Most Luminous Hot Dog



# Summary

- WISE is a great tool for finding AGN, particularly obscured ones.
- WISE can help us learn about the properties of the “normal” AGN torus from “simple” counting statistics.
- WISE has identified some of the most extreme AGN dominated galaxies in the Universe
  - Hot DOGs have unique properties, unlike those of any other luminous galaxy population
  - Possibly a significant stage in the evolution of massive galaxies.
- **Many thanks to the entire WISE team!**



# Backup Slides

# Spitzer AGN Selection

Stern et al. (2005, ApJ, 631, 163)

80% reliable, 80% complete,  $[4.5] \approx 16$

70% reliable, 50% complete,  $[4.5] \approx 17$





Lacy et al. (2004, ApJS, 154, 166)

40% reliable, 90% complete,  $[4.5] \approx 16$

40% reliable, 80% complete,  $[4.5] \approx 17$


# UV – Mid-IR SEDs






Adapted from Eisenhardt et al.(2012)

# AGN Obscuration



- 15-30 mags of extinction
  - Assuming  $R_v = 3.1$
  - Compton thick?
- W1814+3412 more obscured than typical

# Broad Halpha + Outflows?




# The Role of WISE in Finding AGN

The mid-IR is great for

- Particularly at 3-5  $\mu$ m
- Hot dust from the Torus is
- Most stars are red (Raylei

We have known this:

- Lacy et al. (2004, 2005)
- Stern et al. (2005)
- Alonso-Herrero et al. (2005)
- Donley et al. (2008, 2009)
- Assef et al. (2010)



# Completeness/Reliability Fraction

